

# Motion in Two and Three Dimensions: Vectors


Physics 1425 Lecture 4

# Today's Topics

- In the previous lecture, we analyzed the motion of a particle moving vertically under gravity.
- In this lecture and the next, we'll generalize to the case of a particle moving in two or three dimensions under gravity, like a **projectile**.
- **First** we must generalize displacement, velocity and acceleration to two and three dimensions: these generalizations are **vectors**.

# Displacement


- We'll work usually in two dimensions—the three dimensional description is very similar.
- Suppose we move a ball from point **A** to point **B** on a tabletop. This **displacement** can be fully described by giving a **distance** and a **direction**.
- Both can be represented by an arrow, the length some agreed scale: arrow length 10 cm representing 1 m displacement, say.
- This is a **vector**, written with an arrow  $\vec{r}$ : it has **magnitude**, meaning its length, written  $|\vec{r}|$ , and direction.



# Displacement as a Vector

- Now move the ball a *second* time. It is evident that the total displacement, the sum of the two, called the **resultant**, is given by adding the two vectors tip to tail as shown:

- Adding displacement vectors (and notation!):




# Adding Vectors

- You can see that

$$\vec{r}_1 + \vec{r}_2 = \vec{r}_2 + \vec{r}_1.$$

- The vector  $\vec{r}_1$  represents a **displacement**, like saying walk 3 meters in a north-east direction: **it works from any starting point**.


- Adding vectors :



# Subtracting Vectors

- It's pretty easy: just ask, what vector has to be added to  $\vec{a}$  to get  $\vec{b}$  ?
- The answer must be  $\vec{b} - \vec{a}$
- To construct it, put the **tails** of  $\vec{a}, \vec{b}$  together, and draw the vector from the head of  $\vec{a}$  to the head of  $\vec{b}$ .

*Finding the difference:*



# Multiplying Vectors by Numbers

- Only the **length** changes: the direction stays the same.




- Multiplying and adding or subtracting:



# Vector Components

- Vectors can be related to the more familiar Cartesian coordinates  $(x, y)$  of a point  $P$  in a plane: suppose  $P$  is reached from the origin by a displacement  $\vec{r}$ .
- Then  $\vec{r}$  can be written as the **sum of successive displacements in the  $x$ - and  $y$ -directions**:
- These are called the **components** of  $\vec{r}$ .
- Define  $\hat{i}, \hat{j}$  to be vectors of **unit length** parallel to the  $x, y$  axes respectively. The components are  $x\hat{i}, y\hat{j}$ .






# How $\vec{r}$ Relates to $(x, y)$

- The length (magnitude) of  $\vec{r}$  is

$$|\vec{r}| = \sqrt{x^2 + y^2}$$

The angle between the vector and the x-axis is given by:

$$\tan \theta = \frac{y}{x}.$$




# Average Velocity in Two Dimensions

average velocity = displacement/time

- A

In moving from point  $\vec{r}_1$  to  $\vec{r}_2$ , the **average velocity** is in the direction  $\vec{r}_2 - \vec{r}_1$ :

$$\vec{v} = \frac{\vec{r}_2 - \vec{r}_1}{t_2 - t_1}$$




# Instantaneous Velocity in Two Dimensions

$$\vec{v} = \lim_{\Delta t \rightarrow 0} \frac{\Delta \vec{r}}{\Delta t} = \frac{d\vec{r}}{dt}$$


- Note:  $\Delta \vec{r}$  is small, but that doesn't mean  $\vec{v}$  has to be small—  $\Delta t$  is small too!

Defined as the average velocity over a vanishingly small time interval : points in direction of motion at that instant:




# Average Acceleration in Two Dimensions

- Car moving along curving road:




$$\vec{a} = \frac{\vec{v}_2 - \vec{v}_1}{t_2 - t_1}$$



Note that the velocity vectors **tails** must be together to find the difference between them.

# Instantaneous Acceleration in Two Dimensions

$$\vec{a} = \lim_{\Delta t \rightarrow 0} \frac{\Delta \vec{v}}{\Delta t} = \frac{d\vec{v}}{dt}$$



# Acceleration in Vector Components

$$\vec{a} = \frac{d\vec{v}}{dt} = \frac{d}{dt} \left( \frac{d\vec{r}}{dt} \right) = \frac{d^2\vec{r}}{dt^2}$$

Writing  $\vec{a} = (a_x, a_y)$ ,  $\vec{r} = (x, y)$  and matching:

$$a_x = \frac{d^2x}{dt^2}, \quad a_y = \frac{d^2y}{dt^2}$$

as you would expect from the one-dimensional case.

## Clicker Question

A car is moving around a circular track at a constant speed. What can you say about its acceleration?

- A. It's along the track
- B. It's outwards, away from the center of the circle
- C. It's inwards
- D. There is no acceleration

# Relative Velocity

## Running Across a Ship

- A cruise ship is going north at 4 m/s through still water.
- You jog at 3 m/s directly across the ship from one side to the other.
- What is your velocity *relative to the water*?






# Relative Velocities **Just Add...**

- If the **ship's velocity** relative to the water is  $\vec{v}_1$
- And **your velocity** relative to the ship is  $\vec{v}_2$
- Then **your velocity** relative to the water is

$$\vec{v}_1 + \vec{v}_2$$



- Hint: think how far you are *displaced* in one second!